

SLOAN/JOHNSON-COUSINS/2MASS COLOR TRANSFORMATIONS FOR COOL-STARs

JAMES R. A. DAVENPORT

Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195

ANDREW A. WEST

Astronomy Department, University of California, 601 Campbell Hall, Berkeley, CA 94720-3411

CALEB K. MATTHIESEN

Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195

MICHAEL SCHMIEDING

Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195

ADAM KOBELSKI

Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195 and
 Department of Physics, 264 EPS Building, Montana State University, Bozeman, MT 59717

Accepted for Publication in PASP

ABSTRACT

We present multi-color transformations and photometric parallaxes for a sample of 40 low mass dwarfs selected from the Sloan Digital Sky Survey (SDSS) and the General Catalog of Trigonometric Stellar Parallaxes. Our sample was re-observed at the Manastash Ridge Observatory (MRO) using both Sloan and Johnson-Cousins filters and color transformations between the two photometric systems were derived. A subset of the sample had previously measured Johnson-Cousins photometry and parallaxes as well as 2MASS photometry. We observed these stars at MRO using Sloan filters and used these data to derive photometric parallax relations as well as SDSS/Johnson-Cousins/2MASS color transformations. We present the data and derived transformations for use in future low mass star studies.

Subject headings: solar neighborhood — stars: low-mass, brown dwarfs — stars: late-type — stars: distances

1. INTRODUCTION

The most numerous stars in the Galaxy are low-mass dwarfs and yet it is only in the past couple of decades that large enough samples existed to statistically study their properties. The Palomar/Michigan State University (PMSU) survey of low-mass stars (Reid et al. 1995; Hawley et al. 1996; Gizis et al. 2002; Reid et al. 2002;) examined the photometric, spectroscopic and dynamical properties of low mass dwarfs with unprecedented statistical accuracy. Recently, Bochanski et al. (2005) presented another statistical study of the spectroscopic/kinematic properties of nearby low mass dwarfs within 100 pc of the Sun. The advent of large surveys such as the Sloan Digital Sky Survey (SDSS; Gunn et al. 1998; Fukugita et al. 1996; Lupton et al. 1999; York et al. 2000; Hogg et al. 2001; Gunn et al. 2006; Ivezić et al. 2004; Pier et al. 2003; Smith et al. 2002; Stoughton et al. 2002) has increased the sample size and statistical robustness of low-mass star studies (Hawley et al. 2002; West et al. 2004; Silvestri et al. 2006; West et al. 2006; Bochanski et al. 2006; Hawley et al. in prep).

One of the challenges with the SDSS samples has been the ability to accurately compare the Sloan photometry to the Johnson-Cousins standards for low-mass stars

that are used in other studies. Smith et al. (2002) previously investigated the stellar SDSS/Johnson-Cousins color transformation using the 158 SDSS standard stars. However, the SDSS standard stars only extend to an $r - i < 1.5$ (spectral type $\sim$ M4). Recently, Jordi, Grebel & Ammon (2006) and Rodgers et al. (2006) examined a total of several hundred stars and derived new SDSS/Johnson-Cousins transformations. Unfortunately, neither of these new sets of relations extend far beyond $r - i \sim 1.5$ and therefore are not useful for cool-star studies.

In another study, Walkowicz, Hawley & West (2004) projected Sloan filter curves onto spectra of stars with measured Johnson-Cousins photometry in order to derive color transformations between the two systems. Although the Walkowicz et al. method has its utility and has been used by several subsequent studies (West et al. 2004; West, Walkowicz & Hawley 2005; Silvestri et al. 2006), it is not the ideal method for deriving color transformations.

Walkowicz et al. (2004) also derived a relation between Sloan colors and absolute magnitude for stars with measured parallaxes. However, this relation used the intermediate step of estimating the Sloan colors from either the spectral type of the star or the aforementioned transformations. No stars with known parallaxes were

observed directly with both filter sets.

In this paper, we undertake the more ideal solution of observing a set of low-mass dwarfs with the same telescope in both Johnson-Cousins and Sloan filters and using these data to derive a new set of color transformations. A subset of our stars have measured parallaxes and 2MASS photometry and we use these data to derive SDSS/Johnson-Cousins/2MASS color transformations as well as new photometric parallax relations for all three systems. In §2, we describe our sample selection. We detail our observations in §3 and present our results in §4. We discuss these our results and possibilities for the future in §5.

2. SAMPLE SELECTION

2.1. SDSS Selected Sample

The SDSS/Johnson-Cousins color transformation sub-sample was selected from the SDSS Data Release 4 (DR4; Abazajian et al. 2005) database. We required that stars have colors $0.25 < r - i < 3.5$ and $0.75 < g - r < 2.0$ based on the colors for cool-stars reported in West et al. (2004, 2005) and Covey et al. (in prep). In addition, we selected stars with $15 < r < 18.5$, and to ensure a clean sample, made sure that the SDSS flags SATURATED, EDGE, DEBLENDED_AS_MOVING, CHILD, INTERP_CENTER, PSF_FLUX_INTERP and BLENDED were unset. We selected the 5 brightest, observable (based on RA) stars from every 0.1 bin in $r - i$ color (if 5 existed), resulting in a sample of 23 stars. The sample was then divided into a “blue” sample ($r - i < 1.4$; 11 stars) and a “red” sample ($r - i > 1.4$; 12 stars).

2.2. Parallax Selected Sample

The parallax sub-sample is composed of 17 stars selected from the General Catalogue of Trigonometric Stellar Parallaxes (van Altena, Lee & Hoffleit 1995). We first queried the catalog for those stars with small parallax errors ($\sigma \lesssim 0.01''$) and late spectral types (K0 and later). The final sub-sample was selected to span as large a range of spectral types as possible, while maximizing observational efficiency (including both location on the sky and stellar magnitude).

2MASS photometry was obtained from the 2MASS point source catalog (Skrutskie et al. 2006) and the Johnson-Cousin colors were taken from the PMSU I (Reid et al. 1995).

3. OBSERVATIONS

Observations were carried out during the summer of 2005 using the University of Washington 30-inch telescope at Manastash Ridge Observatory (MRO). We used the 2048×2048 CCD detector and the MRO suite of $VRiriz$ filters. Standard stars were observed to derive extinction coefficients and photometric zero-points. Every sample and standard star was observed long enough to obtain a minimum of 10000 counts, and therefore a 1% photon noise uncertainty. The MRO photometric camera was shown to be linear over the entire magnitude range of our observations. The data were reduced using the standard IRAF packages. All Sloan bands were reduced to the SDSS 2.5m system (unprimed). The uncertainty in the derived extinction coefficients dominates the reported photometric uncertainties.

For the SDSS selected sample, we observed each star using both the Johnson-Cousins RI filters and the Sloan ri filters. For the blue sample, we added a Johnson-Cousins V filter and for the red sample the z filter was included. Because the stars in the SDSS selected sample already had SDSS photometry, the MRO riz data allow us ensure that our photometry is correctly on the proper SDSS system. We find that the $r - i$ colors have a mean difference of 0.054 mags and an rms scatter of 0.165 mags, while the $i - z$ colors have a mean offset of -0.002 mags and an rms scatter of 0.220 mags.

4. RESULTS

The resulting photometry can be found in Tables 1 and 2. The observational uncertainties reported in Tables 1 and 2 reflect the propagated uncertainties from each stage of the photometric reductions. Using these new colors we derived new color-color transformations between the Sloan, Johnson-Cousins and 2MASS photometric systems.

4.1. Color-Color Transformations

We performed polynomial fits for each color-color relation and derive the color transformation equations shown below. Each equation contains the uncertainty in the resulting transformed color. The equations and uncertainties derived in this section are only valid over the range in which we have data. These relations should not be extrapolated to redder or bluer colors.

The relations for $R - I$ and $V - R$ as a function of $r - i$ are shown in Figure 1 and Figure 2, are given by:

$$R - I = 0.14 + 0.38(r - i) + 0.29(r - i)^2 + 0.06(r - i)^3 \pm 0.12 \quad (1)$$

and

$$V - R = 0.17 + 1.63(r - i) - 1.19(r - i)^2 + 0.37(r - i)^3 \pm 0.06 \quad (2)$$

and are valid over the range $0.33 \leq r - i \leq 2.85$. The $R - I$ vs. $r - i$ relations of Jordi et al. (2006) and Rodgers et al. (2006) have been overplotted (dotted and dashed respectively) on Figure 1. The three relations agree within the uncertainties over the overlapping region.

The transformation between $R - I$ and $i - z$ is shown in Figure 3, described by:

$$R - I = 0.01 + 2.98(i - z) - 1.51(i - z)^2 + 0.46(i - z)^3 \pm 0.17 \quad (3)$$

and is valid over the range $0.17 \leq i - z \leq 1.59$.

Although the uncertainties in J are large, for completeness we included a first-order fit to the $R - I$ vs $i - J$ relation as Figure 4 and given by:

$$R - I = -0.35 + 0.85(i - J) \pm 0.13 \quad (4)$$

This relation is valid over the range $1.37 \leq i - J \leq 2.58$.




FIG. 1.— $R-I$ vs. $r-i$ color transformation for low-mass dwarfs. The derived polynomial relation is overplotted (solid) and is given by Equation 1. The relations from Jordi et al. (2006) and Rodgers et al. (2006) are overplotted for comparison (dotted and dashed respectively). The three observationally derived relations agree to within the uncertainties.




FIG. 2.— $V-R$ vs. $r-i$ color transformation for the bluest stars in our sample. The derived polynomial relation is overplotted and is given by Equation 2.




FIG. 3.— $R-I$ vs. $i-z$ color transformation for low-mass dwarfs. The derived polynomial relation is overplotted and is given by Equation 3.




FIG. 4.— $R-I$ vs. $i-J$ color transformation for stars with 2MASS data. The derived linear relation is overplotted and is given by Equation 4.




FIG. 5.— $r-i$ vs. $V-I$ color transformation for the bluest stars in our sample. The derived polynomial relation is overplotted (solid) and is given by Equation 5. The spectroscopically derived relation of West et al. (2005) is also overplotted (dashed).

We derived the color-color transformations for $r-i$ vs. $V-I$ to compare with the same relation derived by West et al. (2005; hereafter W05). Although our sample does not go as red and is slightly bluer than the W05. sample, the overlap region allows for useful comparisons.

Figure 5 shows the $r-i$ vs. $V-I$ with the W05 transformation overplotted (dashed line). Our new relation is given by:

$$r-i = -0.76 + 1.73(V-I) - 0.85(V-I)^2 + 0.19(V-I)^3 \pm 0.07 \quad (5)$$

for the ranges $0.95 \leq V-I \leq 2.59$. Our relation agrees with that of W05, confirming the accuracy and feasibility of the W05 method and extending the existing transformations to bluer colors.

4.2. Photometric Parallax




FIG. 6.— Absolute i -band magnitude as a function of $i - z$ color for the stars with direct trigonometric parallax measurements. The best-fit photometric parallax relation is plotted (solid) and given by Equation 6. The West et al. (2005) relation is overplotted (dashed) for comparison.

Using the stars with direct parallax measurements we derive photometric parallax relations for the $i - z$ and $i - J$ colors. These relations are the same relations derived by W05. Although some of the uncertainties in the stellar photometry are large, these data serve as a direct confirmation of the W05 relations, without having to transform to a different photometric system.

The relations are shown in Figures 6 and 7. The fits are plotted as solid lines and given by the equations,

$$M_i = 5.49 + 6.63(i - z) \pm 0.44 \quad (6)$$

and

$$M_i = 2.55 + 3.38(i - J) \pm 0.63 \quad (7)$$

and are valid over the ranges $0.22 \leq i - z \leq 0.98$ and $1.37 \leq i - J \leq 2.58$ respectively. The W05 photometric parallax relations are shown as dashed lines. Both of the fits from this study are consistent with the W05 results and extend the relations to bluer colors. The $i - J$ relation of W05 can be extended to bluer colors and appears to follow the same relation to within the uncertainties. Although the piece-wise nature of the W05 $i - z$ relation precludes an extrapolation to bluer colors, the $i - z$ relation from this paper provides a useful extension for bluer stars.

5. DISCUSSION

We have derived SDSS-Johnson-2MASS color transformations based on direct measurements of cool-stars. We present several relations not included in previous studies of low-mass stars. Our $r - i$ vs. $V - I$ relation agrees with the previous findings of W05. In addition, we extend many of the relations to bluer colors than those included in the W05 analysis and redder colors than the Smith et al. (2002), Rodgers et al. (2006) and Jordi et al. (2006). Table 3 summarizes our derived color transformations and compares them with the relations given in W05, Rodgers et al. (2006) and Jordi et al. (2006). This table is currently the most complete compilation of low-mass star color transformations.




FIG. 7.— Absolute i -band magnitude as a function of $i - J$ color for the stars with direct trigonometric parallax measurements. The best-fit photometric parallax relation is plotted (solid) and given by Equation 7. The West et al. (2005) relation is overplotted (dashed) for comparison.

We also utilized the stars in our sample that have direct parallax measurements to derive photometric parallax relations. Our results confirm the relations given by W05.

All of our results indicate that the W05 method is justified and that the spectroscopically derived W05 color transformations and spectroscopic parallax relations are valid to within the quoted uncertainties. Future studies with larger samples of cool-stars will further constrain these relations.

The authors would like to thank Kevin Covey for his useful discussions and many suggestions while working on this paper. We would also like to thank Cindy Lee for helping with the observations, Suzanne Hawley for her helpful suggestions and Chris Laws for his support and care of the MRO facilities. AAW acknowledges the support of NSF grant AST05-40567 and the financial support of Julianne Dalcanton.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, Cambridge University, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-

Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton Uni-

versity, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Abazajian, K., et al. 2004, *AJ*, 128, 502
 Adelman-McCarthy, J. K., et al. 2006, *ApJS*, 162, 38
 Bochanski, J. J., Hawley, S. L., Reid, I. N., Covey, K. R., West, A. A., Tinney, C. G., & Gizis, J. E. 2005, *AJ*, 130, 1871
 Bochanski, J. J., West, A. A., Hawley, S. L., & Covey, K. R. 2006, *AJ*, in press
 Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
 Gizis, J. E., Reid, I. N., & Hawley, S. L. 2002, *AJ*, 123, 3356
 Gunn, J. E., et al. 1998, *AJ*, 116, 3040
 —. 2006, *AJ*, 131, 2332
 Hawley, S. L., Covey, K. R., et al. 2002, *AJ*, 123, 3409
 Hawley, S. L., Gizis, J. E., & Reid, I. N. 1996, *AJ*, 112, 2799
 Hogg, D. W., Finkbeiner, D. P., Schlegel, D. J., & Gunn, J. E. 2001, *AJ*, 122, 2129
 Ivezić, Ž., et al. 2004, *Astronomische Nachrichten*, 325, 583
 Jordi, K., Grebel E. K., & Ammon, K. 2006, *A&A*, in press
 Lupton, R. H., Gunn, J. E., & Szalay, A. S. 1999, *AJ*, 118, 1406
 Pier, J. R., Munn, J. A., Hindsley, R. B., Hennessy, G. S., Kent, S. M., Lupton, R. H., & Ivezić, Ž. 2003, *AJ*, 125, 1559
 Reid, I. N., Gizis, J. E., & Hawley, S. L. 2002, *AJ*, 124, 2721
 Reid, I. N., Hawley, S. L., & Gizis, J. E. 1995, *AJ*, 110, 1838
 Rodgers, C. T., Canterna, R., Smith, J. A., Pierce, M. J., & Tucker, D. L. 2006, *AJ*, 132, 989
 Silvestri, N. M., et al. 2006, *AJ*, 131, 1674
 Skrutskie, M. F., et al. 2006, *AJ*, 131, 1163
 Smith, J. A., et al. 2002, *AJ*, 123, 2121
 Stoughton, C., et al. 2002, *AJ*, 123, 485
 Strauss, M. A., et al. 2002, *AJ*, 124, 1810
 van Altena, W. F., Lee, J. T., & Hoffleit, E. D. 1995, *The general catalogue of trigonometric [stellar] parallaxes* (New Haven, CT: Yale University Observatory, —c1995, 4th ed., completely revised and enlarged)
 Walkowicz, L. M., Hawley, S. L., & West, A. A. 2004, *PASP*, 116, 1105
 West, A. A., Hawley, S. L., Walkowicz, L. M., Covey, K. R., Silvestri, N. M., Raymond, S. N., Harris, H. C., Munn, J. A., McGehee, P. M., Ivezić, Ž., & Brinkmann, J. 2004, *AJ*, 128, 426
 West, A. A., Walkowicz, L. M., & Hawley, S. L. 2005, *PASP*, 117, 706
 West, A. A., Bochanski, J. J., Hawley, S. L., Cruz, K. L., Covey, K. R., Silvestri, N. M., Reid, I. N., & Liebert, J. 2006, *AJ*, in press
 York, D. G., et al. 2000, *AJ*, 120, 1579

TABLE 1
PHOTOMETRY FOR SDSS SELECTED STARS

RA (J2000)	DEC (J2000)	$V - R$	R	$R - I$	r	$r - i$	$i - z$
12:34:57.5	+31:12:22	0.94 (0.03)	13.99 (0.71)	1.01 (0.12)	15.02 (0.02)	0.73 (0.02)	0.43 (0.02)
13:26:16.3	+56:40:44		13.44 (0.35)	2.94 (0.08)	17.68 (0.01)	2.52 (0.02)	1.40 (0.02)
13:49:06.4	+60:22:20	0.76 (0.03)	15.61 (0.71)	0.48 (0.12)	15.00 (0.02)	0.43 (0.03)	0.23 (0.03)
14:09:12.8	+65:00:00		15.49 (0.62)	1.61 (0.09)	15.91 (0.02)	1.60 (0.02)	0.86(0.02)
14:30:57.4	+51:24:52	0.80 (0.02)	15.38 (0.71)	0.62 (0.12)	15.01 (0.01)	0.53 (0.02)	0.28 (0.02)
14:44:17.1	+30:02:12		16.06 (0.62)	2.64 (0.11)	18.24 (0.02)	2.85 (0.03)	1.59 (0.03)
14:44:26.2	+62:01:25	0.76 (0.04)	15.35 (0.36)	0.56 (0.08)	15.00 (0.02)	0.53 (0.02)	0.30 (0.02)
15:06:24.5	+16:27:53	0.72 (0.04)	15.26 (0.36)	0.51 (0.08)	15.00 (0.02)	0.46 (0.03)	0.28 (0.02)
15:23:42.4	+35:47:09		14.30 (0.62)	1.18 (0.09)	15.06 (0.02)	0.93 (0.02)	0.49 (0.02)
15:27:07.4	+43:03:01		14.36 (0.62)	1.35 (0.09)	15.24 (0.01)	1.10 (0.02)	0.57 (0.02)
15:34:41.4	+46:41:32		17.06 (0.35)	2.03 (0.09)	17.43 (0.01)	2.27 (0.02)	1.24 (0.02)
15:46:05.3	+37:49:45		17.92 (0.37)	2.62 (0.14)	18.44 (0.02)	2.52 (0.02)	1.43 (0.02)
16:00:15.6	+52:52:27	0.56 (0.04)	15.39 (0.36)	0.39 (0.08)	15.00 (0.01)	0.33 (0.02)	0.16 (0.03)
16:11:39.8	+40:05:59		16.77 (0.35)	1.72 (0.09)	16.94 (0.02)	1.83 (0.02)	0.95 (0.02)
16:27:18.2	+35:38:35		15.93 (0.36)	2.36 (0.10)	17.72 (0.02)	2.29 (0.02)	1.34 (0.03)
16:37:18.6	+44:28:46		15.24 (0.61)	1.60 (0.09)	15.72 (0.01)	1.55 (0.02)	0.83 (0.02)
17:18:18.3	+65:12:12	0.56 (0.04)	15.34 (0.36)	0.44 (0.08)	15.00 (0.01)	0.29 (0.02)	0.17 (0.02)
17:30:17.3	+62:29:26	1.23 (0.03)	15.80 (0.71)	1.36 (0.12)	15.45 (0.02)	1.39 (0.02)	0.71 (0.03)
21:36:32.5	-06:41:51	0.57 (0.05)	13.89 (0.36)	0.70 (0.08)	15.02 (0.01)	0.38 (0.02)	0.17 (0.02)
21:37:27.5	-08:27:13	0.78 (0.06)	13.86 (0.36)	0.88 (0.08)	15.02 (0.01)	0.77 (0.02)	0.44 (0.02)
23:21:24.0	-09:57:36	0.96 (0.04)	14.32 (0.71)	1.15 (0.12)	15.01 (0.02)	0.94 (0.03)	0.50 (0.02)
23:46:44.7	+16:03:59		16.47 (0.62)	2.24 (0.09)	17.38 (0.01)	2.23 (0.02)	1.36 (0.03)
23:50:47.8	+14:42:44		15.21 (0.35)	1.67 (0.09)	16.39 (0.01)	1.50 (0.02)	0.79 (0.02)

NOTE. — The Sloan photometry comes from SDSS DR4 database. The Johnson/Cousins magnitudes were measured from MRO data. Photometric uncertainties are included in parentheses.

TABLE 2
PHOTOMETRY FOR PARALLAX SELECTED SAMPLE

RA (J2000)	DEC (J2000)	MRO ^a /2MASS ^b					parallax ^c ($''$)	PMSU ^d		
		$r - i$	i	$i - z$	J	$i - J$		V	R	I
12:49:02.7	+66:06:37	1.30 (0.14)	8.91 (0.08)	0.54 (0.14)	6.88 (0.02)	2.03 (0.09)	0.1020 (0.0047)	10.87	9.75	8.28
13:19:45.7	+47:46:39	0.85 (0.03)	7.02 (0.14)	0.22 (0.03)	5.34 (0.03)	1.68 (0.14)	0.1050 (0.0061)	9.07	8.16	7.18
14:02:33.3	+46:20:24	0.90 (0.02)	7.79 (0.11)	0.43 (0.04)	6.26 (0.09)	1.53 (0.14)	0.0890 (0.0039)	...	...	...
14:53:51.2	+23:33:21	1.38 (0.05)	9.71 (0.27)	0.67 (0.06)	7.44 (0.02)	2.27 (0.27)	0.0960 (0.0041)	...	...	...
16:02:50.6	+20:35:16	1.47 (0.06)	10.57 (0.29)	0.72 (0.06)	8.13 (0.02)	2.44 (0.29)	0.1000 (0.0031)	...	...	...
16:24:09.5	+48:21:10	1.13 (0.04)	8.55 (0.13)	0.59 (0.02)	6.64 (0.02)	1.91 (0.13)	0.1310 (0.0048)	10.73	9.74	8.43
17:07:07.4	+21:33:14	1.34 (0.03)	9.78 (0.09)	0.56 (0.01)	7.88 (0.02)	1.90 (0.09)	0.0680 (0.0029)	10.96	9.91	8.55
17:19:54.2	+26:30:04	1.31 (0.13)	9.30 (0.08)	0.51 (0.13)	7.27 (0.02)	2.03 (0.08)	0.0940 (0.0021)	11.03	9.89	8.36
18:35:20.3	+45:45:06	0.72 (0.02)	8.49 (0.24)	0.39 (0.04)	6.88 (0.02)	1.61 (0.24)	0.0670 (0.0020)	...	...	...
19:21:34.0	+20:50:24	1.57 (0.14)	11.27 (0.09)	0.77 (0.15)	8.80 (0.02)	2.47 (0.09)	0.1000 (0.0030)	...	...	...
20:03:24.8	+29:52:00	1.71 (0.07)	12.10 (0.33)	0.91 (0.06)	9.55 (0.02)	2.55 (0.33)	0.0860 (0.0228)	13.34	12.01	10.23
20:05:02.8	+54:25:43	0.81 (0.04)	10.49 (0.29)	0.47 (0.06)	8.93 (0.02)	1.56 (0.29)	0.0610 (0.0022)	10.98	10.03	8.99
20:29:53.1	+09:40:17	1.73 (0.15)	10.90 (0.10)	0.87 (0.16)	8.23 (0.02)	2.67 (0.10)	0.1140 (0.0019)	...	...	...
20:40:40.4	+15:29:51	1.66 (0.06)	11.13 (0.31)	0.86 (0.07)	8.64 (0.03)	2.49 (0.31)	0.1020 (0.0102)	...	...	...
20:41:28.4	+57:25:08	0.61 (0.10)	9.01 (0.06)	0.28 (0.09)	7.64 (0.02)	1.37 (0.06)	0.0440 (0.0102)	...	...	...
20:43:25.8	+55:22:55	1.88 (0.07)	12.14 (0.33)	0.98 (0.08)	9.56 (0.02)	2.58 (0.33)	0.0630 (0.0055)	...	...	...
23:43:11.8	+36:32:08	1.72 (0.14)	10.34 (0.09)	0.71 (0.15)	8.11 (0.03)	2.23 (0.10)	0.1230 (0.0029)	...	...	...

NOTE. — Uncertainties are included in parentheses where available.

^a Sloan magnitudes and colors derived from MRO photometric data.

^b 2MASS point source catalog photometry (Skrutskie et al. 2006).

^c Parallaxes from General Catalogue of Trigonometric Stellar Parallaxes (van Altena et al. 1995).

^d Johnson-Cousins photometry from PMSU I (Reid et al. 1995).

TABLE 3
SUMMARY OF COLOR TRANSFORMATIONS

Davenport et al. ^a	West et al. (2005) ^b	Rodgers et al. (2006) ^c	Jordi et al. (2006) ^d
$V - R = 0.17 + 1.63(r - i) - 1.19(r - i)^2$ $+0.37(r - i)^3 \pm 0.06$	...	...	$g - r = (1.646 \pm 0.008)(V - R)$ -0.139 ± 0.004
$R - I = 0.14 + 0.38(r - i) + 0.29(r - i)^2$ $+0.06(r - i)^3 \pm 0.12$	...	$r - i = (1.000 \pm 0.006)(R - I) - 0.212$	$r - i = (1.007 \pm 0.005)(R - I)$ -0.236 ± 0.003
$R - I = 0.01 + 2.98(i - z) - 1.51(i - z)^2$ $+0.46(i - z)^3 \pm 0.17$	...	$r - z = (1.567 \pm 0.020)(R - I) - 0.365$	$r - z = (1.584 \pm 0.008)(R - I)$ -0.386 ± 0.005
$R - I = -0.35 + 0.85(i - J) \pm 0.13$	$i - z = -20.6 + 26.0(I - K) - 11.7(I - K)^2$ $+2.30(I - K)^3 - 0.17(I - K)^4$	...	...
$r - i = -0.76 + 1.73(V - I) - 0.85(V - I)^2$ $+0.19(V - I)^3$	$r - i = -2.69 + 2.29(V - I) - 0.28(V - I)^2$	...	...

^aRelations are valid over $0.17 \leq i - z \leq 1.59$, $0.33 \leq r - i \leq 2.85$, $1.37 \leq i - J \leq 2.58$, and $0.95 \leq V - I \leq 2.59$

^bRelations are valid over $0.67 \leq r - i \leq 2.01$, and $0.37 \leq i - z \leq 1.84$

^cRelations are valid over $-0.095 \leq R - I \leq 0.695$

^dRelations are valid over $0.0 \leq V - R \leq 1.2$, and $0.0 \leq R - I \leq 2.0$